

Bounds on long-lived charged massive particles from Big Bang nucleosynthesis

Karsten Jedamzik

*Laboratoire de Physique Mathématique et Théorique, C.N.R.S.,
Université de Montpellier II, 34095 Montpellier Cedex 5, France*

The Big Bang nucleosynthesis (BBN) in the presence of charged massive particles (CHAMPs) is studied in detail. All currently known effects due to the existence of bound states between CHAMPs and nuclei, including possible late-time destruction of ${}^6\text{Li}$ and ${}^7\text{Li}$ are included. The study sets conservative bounds on CHAMP abundances in the decay time range $3 \times 10^2 \text{sec} \lesssim \tau_x \lesssim 10^{12} \text{sec}$. It is stressed that the production of ${}^6\text{Li}$ at early times $T \sim 10 \text{keV}$ is overestimated by a factor ~ 10 when the approximation of the Saha equation for the ${}^4\text{He}$ bound state fraction is utilised. To obtain conservative limits on the abundance of CHAMPs, a Monte-Carlo analysis with $\sim 3 \times 10^6$ independent BBN runs, varying reaction rates of nineteen different reactions, is performed (see attached erratum, however). The analysis yields the surprising result that except for small areas in the particle parameter space conservative constraints on the abundance of decaying charged particles are currently very close to those of neutral particles. It is shown that, in case the rates of a number of heretofore unconsidered reactions may be determined reliably in future, it is conceivable that the limit on CHAMPs in the early Universe could be tightened by orders of magnitude. An **ERRATUM** gives limits on primordial CHAMP densities when the by Ref. [20] recently more accurately determined CHAMP reaction rates are employed.

I. INTRODUCTION

Big Bang nucleosynthesis (BBN) has proven itself as a powerful tool in constraining the conditions of the early Universe and physics beyond the standard model. Thus bounds on a variety of hypothesis have been derived including, for example, modifications of gravity, baryon inhomogeneity, matter-antimatter domains, non-zero lepton chemical potentials, and relic decaying particles. It has been recently realized that BBN may also place bounds on the abundance of charged, weak-scale mass particles existing during and after BBN. Though it had been noted already earlier that negatively charged, weak-scale mass particles (CHAMPs) form bound-states with positively charged nuclei towards the end of BBN [1], it has only recently been put forward that this may impact considerably the light-element yields synthesized during BBN [2, 3, 4].

Here the most important proposed change is due to a catalysing of reactions such as ${}^2\text{H} + {}^4\text{He} \rightarrow {}^6\text{Li} + \gamma$ [2]. Other less important modifications, concerning constraints on CHAMPs, had also been considered [4, 5, 6]. Being of quadrupole (E2) nature the ${}^2\text{H} + {}^4\text{He}$ reaction has a very small rate (S-factor: 10^{-8}MeV barn) thus yielding typically very little ${}^6\text{Li}/{}^1\text{H} \sim 10^{-14}$ in standard BBN. When the helium-nuclei is in a bound state (${}^4\text{He}-X^-$) the above reaction may be replaced by its photonless analogue: ${}^2\text{H} + ({}^4\text{He}-X^-) \rightarrow {}^6\text{Li} + X^-$, with a cross section estimated orders of magnitude larger than that for the standard BBN ${}^6\text{Li}$ -synthesizing process. Initial estimates for this enhancement factor were given at around 6×10^7 [2, 7]. It was argued that since ~ 1 of all X^- are within bound states with ${}^4\text{He}$ at temperatures $T \lesssim 8 \text{keV}$ (cf. Fig. 3) very small abundances of CHAMPs present at $t \approx 10^4 \text{sec}$ in the early Universe could already overproduce the ${}^6\text{Li}$ isotope with respect to observations [8]. Assuming a (too restrictive) ${}^6\text{Li}/{}^1\text{H} \lesssim 2 \times 10^{-11}$ con-

straint, bounds as strong as $n_{X^-}/s \lesssim 2.5 \times 10^{-17}$, the CHAMP-to-entropy ratio, were derived. These bounds were subsequently weakened by one order of magnitude when a more proper evaluation of the rate for the ${}^2\text{H} + ({}^4\text{He}-X^-) \rightarrow {}^6\text{Li} + X^-$ process [9] was derived. Such bounds have now been utilised by a number of groups to constrain, for example, abundances of supersymmetric staus $\tilde{\tau}$ s in the early Universe [2, 7, 9, 10, 11].

Recently, I have shown that there are several changes to the "naive" picture of synthesis of ${}^6\text{Li}$ in the presence of CHAMPs [12]. First, (${}^4\text{He}-X^-$) bound states may be destroyed during the electromagnetic cascades induced by the decay of CHAMPs, rendering ${}^6\text{Li}$ production in some parts of parameter space much less efficient. More importantly, when CHAMPs are still present at times $t \gtrsim 10^6 \text{sec}$ Big Bang nucleosynthesis enters a second phase of Coulomb-unsuppressed reactions on bound states between charge $Z = 1$ nuclei and a CHAMP. It has been shown that reactions such as ${}^6\text{Li} + ({}^1\text{H}-X^-) \rightarrow {}^4\text{He} + {}^3\text{He} + X^-$, ${}^7\text{Li} + ({}^1\text{H}-X^-) \rightarrow {}^8\text{B} + X^-$, etc, are capable of completely destroying any priorly synthesized ${}^6\text{Li}$ and ${}^7\text{Li}$. This is possible in particular at somewhat higher CHAMP-to-entropy $Y_X = n_X/s$ ratios. In order to estimate the efficiency of such destruction, not only was it required to estimate the cross sections for such ${}^6\text{Li}$ and ${}^7\text{Li}$ destroying reactions, but also those of CHAMP exchange reactions such as $({}^1\text{H}-X^-) + {}^2\text{H} \rightarrow {}^1\text{H} + ({}^2\text{H}-X^-)$, capable of significantly reducing the ${}^1\text{H}$ bound state fraction. Altogether nineteen reactions of significant importance for the late-time nucleosynthesis $t \gtrsim 10^6 \text{sec}$ have been identified. The rates for all these reactions were determined in the Born approximation. Concerning details on the BBN with CHAMPs at late times, the importance of particular reactions, and their evaluation, the reader is referred to the original paper [12]. Unfortunately, the Born approximation is not a particular good approximation for determining rates of these nineteen important reactions




FIG. 1: Limits on the primordial CHAMP-to-entropy ratio $Y_x = n_X/s$ (with $n_{X^-}/s = Y_x/2$) for CHAMPs with intermediate life times. Shown are constraint lines for CHAMPs of mass $M_x = 1$ TeV and a variety of hadronic branching ratios $B_h = 10^{-5} - 1$, as labeled in the figure. Solid (red) lines correspond to the conservativelimit ${}^6\text{Li}/{}^7\text{Li} < 0.66$, whereas dashed (blue) lines correspond to ${}^6\text{Li}/{}^7\text{Li} < 0.1$. It is seen that only for CHAMPs with $B_h \lesssim 10^{-2}$ the effects of bound states become important. For smaller decay times τ_x the limits on CHAMP abundances are virtually identical to those on the abundance of neutral relic decaying particles [14].

for CHAMP BBN, leaving significant uncertainty in the BBN yields with late decaying $\tau_x \gtrsim 10^6$ sec CHAMPs.

Given the above, it seems very premature to rule out CHAMPs in the early Universe simply by their production of ${}^6\text{Li}$ at $T \approx 8$ keV. In this letter constraints on the abundances on CHAMPs are derived which take full account of all the above mentioned extra physics priorly neglected. Here constraints will be placed for two different decay time regimes, for $3 \times 10^2 \text{ sec} \lesssim \tau_x \lesssim 5 \times 10^5 \text{ sec}$ in Section 2, where the important rates are relatively well known, and for $5 \times 10^5 \text{ sec} \lesssim \tau_x \lesssim 10^{12} \text{ sec}$ in Section 3, where a Monte-Carlo analysis is employed to derive conservative limits. It will be seen that constraints change by large factors with respect to those priorly given, particularly for long CHAMP decay times.

II. CONSTRAINTS ON CHAMPS WITH INTERMEDIATE LIFE TIMES

In the next two sections constraints from BBN on the existence of CHAMPs in the early Universe are presented. Such constraints get increasingly more uncertain as the life time τ_x of a CHAMP increases. For $\tau_x \lesssim 3 \times 10^2 \text{ sec}$ CHAMPs have almost no impact on BBN beyond those of their injection of electromagnetically and hadronically interacting particles during their decay [13]. Such constraints have already been discussed in detail in the literature and the reader is referred to, for example, Ref. [14] for details. For $3 \times 10^2 \text{ sec} \lesssim \tau_x \lesssim 5 \times 10^5 \text{ sec}$ con-




FIG. 2: Same as Fig. 1, but for $M_x = 100$ GeV.




FIG. 3: Fraction of CHAMPs f_X which are bound to ${}^4\text{He}$ as a function of temperature for (a) the approximation by the Saha equation, thin-dashed (green) and (b) full numerical integration of the rate equation, thin-solid (red). Also shown is the product of f_X with the rate $\langle \sigma v \rangle$ for the ${}^6\text{Li}$ producing reaction ${}^2\text{H}({}^4\text{He}-X^-, X^-){}^6\text{Li}$, simply denoted as “rate” and in arbitrary units, for both cases (a) thick-dashed (green) and (b) thick-solid (red). The figure illustrates that ${}^6\text{Li}$ production at $T \approx 8$ keV due to bound states is overestimated by a factor ~ 10 when the approximation of the Saha equation is utilised.

straints are still fairly reliable and depend mostly on the ${}^2\text{H}({}^4\text{He}-X^-, X^-){}^6\text{Li}$ rate. Since this one has been determined beyond the Born approximation [9] and is most likely known to within a factor three, limits should also be known up to such a factor. It has been found numerically that other reactions, such as ${}^3\text{H}({}^4\text{He}-X^-, X^-){}^7\text{Li}$ and ${}^3\text{He}({}^4\text{He}-X^-, X^-){}^7\text{Be}$ play less of a role in setting constraints at early times, even in case their rates significantly exceed those determined in the Born approximation.

When deriving constraints the following conservative observationally determined limits on the light element

abundances are adopted:

$$\begin{aligned}
 Y_p &< 0.258 \\
 1.2 \times 10^{-5} &< {}^2\text{H}/\text{H} < 5.3 \times 10^{-5} \\
 {}^3\text{He}/{}^2\text{H} &< 1.52 \\
 8.5 \times 10^{-11} &< {}^7\text{Li}/\text{H} < 5 \times 10^{-10} \\
 {}^6\text{Li}/{}^7\text{Li} &< 0.66 \quad (0.1)
 \end{aligned} \tag{1}$$

Here Y_p denotes the helium mass fraction. The observations behind these limits are discussed in further detail in Ref. [14]. It should be noted here that the frequently used bound ${}^6\text{Li}/{}^1\text{H} \lesssim 2 \times 10^{-11}$ is too stringent. ${}^6\text{Li}$ may be destroyed during the life time of a Population II star. In fact, if the current discrepancy between standard BBN predicted ${}^7\text{Li}/{}^1\text{H} \approx 4 - 5 \times 10^{-10}$ and Pop II star observed ${}^7\text{Li}/{}^1\text{H} \approx 1 - 2.5 \times 10^{-10}$ is resolved by factor 2 – 3 stellar ${}^7\text{Li}$ destruction, as claimed for example in Ref. [15], than ${}^6\text{Li}$ is destroyed by at least the same factor. This would yield an upper limit close to ${}^6\text{Li}/{}^1\text{H} \lesssim 4 \times 10^{-11}$, or ${}^6\text{Li}/{}^7\text{Li} \lesssim 0.1$. However, since ${}^6\text{Li}$ is more fragile than ${}^7\text{Li}$ it may, in principle, be destroyed by much larger factors than ${}^7\text{Li}$, as shown in a number of stellar evolution studies [16]. A conservative limit of ${}^6\text{Li}/{}^7\text{Li} \lesssim 0.66$ (corresponding approximately to ${}^6\text{Li} \lesssim 2.7 \times 10^{-10}$) was therefore applied in Ref. [14].

In Fig. 1 and Fig. 2 constraints on decaying CHAMPs with intermediate life times are presented. In both figures results are shown for a variety of hadronic branching ratios B_h , with Fig. 1 showing results for $M_x = 1$ TeV, and Fig. 2 for $M_x = 100$ GeV. In order to derive these constraints the rate for ${}^2\text{H}({}^4\text{He}-X^-, X^-){}^6\text{Li}$ as given in Ref. [9] has been utilised. It is seen that for large hadronic branching ratio ($B_h \gtrsim 0.01$ for $M_x = 1$ TeV and $B_h \gtrsim 0.1$ for $M_x = 100$ GeV) constraints depend almost linearly on B_h and are *not* different from those for neutral particles. This seems somewhat surprising, due to the advocated power [2, 9] of helium-CHAMP bound states to produce ${}^6\text{Li}$. Nevertheless, it is known that ${}^6\text{Li}$ is also produced abundantly by hadronic decays during that time [14, 17], and this ${}^6\text{Li}$ source is more important than that of ${}^2\text{H}({}^4\text{He}-X^-, X^-){}^6\text{Li}$ at large B_h . The efficiency of ${}^2\text{H}({}^4\text{He}-X^-, X^-){}^6\text{Li}$ has in any case been overestimated in some papers [9, 10]. ${}^6\text{Li}$ production here is given by a convolution of two exponentials, (a) the exponentially rising ${}^4\text{He}-X^-$ bound state fraction at $T \sim 10$ keV, and (b) the due to a Coulomb barrier exponentially decreasing reaction rate. This leads to the bulk of the ${}^6\text{Li}$ production in a very narrow temperature interval $8 \text{ keV} \gtrsim T \gtrsim 6 \text{ keV}$. Refs. [9] and [10] assumed the applicability of the Saha equation for the ${}^4\text{He}$ bound state fraction (though earlier studies Refs. [2, 3, 4, 5, 7] did not). Numerical integration shows that the formation of appreciable bound state fractions is slightly delayed when compared to the Saha equation, since the recombination rate is of the same order as the Hubble rate. This may be seen in Fig. 3. This slight difference results than in approximately one order of magnitude less ${}^6\text{Li}$ production, due to the convolution of exponen-




FIG. 4: Likelihood areas in the CHAMP-to-entropy ratio Y_X – CHAMP life time τ_X (in seconds) plane for bound-state BBN to obey the observational constraints on light element abundances Eqs. 1. Results of a Monte-Carlo analysis varying nineteen ill-determined reaction rates randomly (see text for details) which significantly impact BBN yields at late times $\tau_x \gtrsim 10^6$ sec are shown. From bottom to top, the areas show likelihoods: $> 99\%$ lightest shade (yellow), $95\% - 99\%$ (green), $80\% - 95\%$ (purple), $20\% - 80\%$ (red), $5\% - 20\%$ (light-blue), $1\% - 5\%$ (black), and $< 1\%$ (dark-blue), respectively, for CHAMP BBN to respect observational constraints. No effects of electromagnetic- and hadronic- cascades due to the CHAMP decay have been taken into account (cf. Fig. 5). The Monte-Carlo analysis presented in this figure employs $\sim 1.5 \times 10^6$ independent BBN calculations.

tials. Very recently, the same observation has also been made in Ref. [18] utilising the set of Boltzman equations relevant for ${}^6\text{Li}$ production via ${}^4\text{He}-X^-$ bound states as given in Ref. [19]. Conservative limits on CHAMPs in the intermediate decay time interval are thus weaker than initially thought. Here the weakening is due to a lower rate [9], the failure of the Saha equation, and a too restrictive upper limit on the ${}^6\text{Li}$ abundance. It is noted here that the photodisintegration of $({}^4\text{He}-X^-)$ bound states as noted in Ref. [12] is comparatively unimportant at small Y_x , such that further weakening of the limit on CHAMPs does not result. In any case, when $B_h \lesssim 10^{-2}$ bound state induced production of ${}^6\text{Li}$ becomes dominant over hadronic production. Since this is the case for, for example, supersymmetric staus, bound state effects thus still remain very constraining in particular scenarios [2].

III. CONSTRAINTS ON CHAMPs WITH LONG LIFE TIMES

When the life times exceeds $\tau_x \gtrsim 5 \times 10^5$ sec it becomes substantially more difficult to place reliable limits. This is due to a large number of CHAMP-induced reactions becoming important at $T \lesssim 1$ keV, in particular all Coulomb-unsuppressed nuclear reactions shown in Table II and Table III of Ref. [12], as well as the CHAMP ex-




FIG. 5: As Fig. 4 but including the effects of electromagnetic cascades during X -decay, assuming that a fraction $f_{EM} = 1$ of the particles rest mass (taken $m_X = 100$ GeV) is converted into electromagnetically interacting particles. The hadronic branching ratio was set to $B_h = 0$. The dashed line shows the analogous limit for neutral relics.




FIG. 6: Isocontours of ${}^6\text{Li}/{}^7\text{Li} = 0.66$ solid (red), and ${}^6\text{Li}/{}^7\text{Li} = 0.1$ dashed (blue), in the CHAMP-to-entropy Y_X - CHAMP life time τ_X plane for rates determined in the unreliable Born approximation [12]. The present figure is *not* supposed to be utilised for limiting CHAMP abundances. It rather is for illustrative purpose, showing the possible importance of the ${}^4\text{He} + ({}^2\text{H}-X^-) \rightarrow X^- + {}^6\text{Li}$ reaction to further tighten limits on long-lived CHAMPS by orders of magnitude due to late-time ${}^6\text{Li}$ production. A hadronic branching ratio of $B_h = 0$ has been assumed.

change reactions shown in Table IV of that paper. This comprises a total number of nineteen reactions. Though all rates have been determined numerically in the Born approximation in Ref. [12], as the Born approximation is likely to fail badly, results become uncertain. In order to still arrive at a reliable result one is thus forced to perform a Monte-Carlo analysis, varying all ill-determined reaction rates within conservative ranges. This has been done in the present paper. In particular, the Born ap-

proximation values of the rates given in Ref. [12] (shown in Figs. 3 and 4, as well as in Table III and IV of that paper), have been taken as benchmarks. For each reaction a random generator determined a factor f_i with which the benchmark rate was multiplied. These factors were generated with a probability distribution flat in logarithmic space, and between values $1/f_i^{cut} \leq f_i \leq f_i^{cut}$. For the reaction-rate dependent conservatively chosen f_i^{cut} the reader is referred to Table VII of Ref. [12]. For each point in parameter space, i.e. for Y_X and τ_X , this procedure was repeated a 1000 times in order to arrive with one thousand different randomly chosen sets for the 19 ill-determined reaction rates. For each realization of reaction rates an independent BBN calculation was then performed and compared to the observational constraints.

One may wonder if 1000 realizations for the reaction rates are actually sufficient for sampling the, a priori, complicated probability space. After all even only adopting two values for each reaction rate, a large rate and a small rate, already yields $2^{19} \approx 5 \times 10^5$ different possibilities for sets of reaction rates. For a few individual points in Y_X and τ_X , I have therefore generated 10^5 different realizations of rate combinations. Comparison with the results by only a 1000 realizations shows that the simulation with a 1000 realizations may be trusted approximately to the 1% level (i.e. < 10 BBN runs passing or failing observational constraints). This is true since at an individual point in the Y_X - τ_X plane, results mostly only depend on a number $\sim 4 - 6$ of rates, with all other rates being less important. Which rates are most important for the BBN yields, however, depends on the location in the Y_X - τ_X plane. For example, at large Y_X (and τ_X) results depend sensitively on the reactions: ${}^3\text{H}({}^4\text{He}-X^-, X^-){}^7\text{Li}$, ${}^6\text{Li}({}^1\text{H}-X^-, X^-){}^4\text{He} + {}^3\text{He}$, and $({}^1\text{H}-X^-) + {}^2\text{H} \rightarrow {}^1\text{H} + ({}^2\text{H}-X^-)$ (and to a lesser degree ${}^7\text{Be}({}^1\text{H}-X^-, X^-){}^8\text{B}$, ${}^4\text{He}({}^3\text{H}-X^-, X^-){}^7\text{Li}$, and $({}^1\text{H}-X^-) + {}^4\text{He} \rightarrow {}^1\text{H} + ({}^4\text{He}-X^-)$), whereas for small Y_X (still large τ_X) they depend mostly on reactions: ${}^4\text{He}({}^2\text{H}-X^-, X^-){}^6\text{Li}$, $({}^1\text{H}-X^-) + {}^2\text{H} \rightarrow {}^1\text{H} + ({}^2\text{H}-X^-)$ and $({}^1\text{H}-X^-) + {}^4\text{He} \rightarrow {}^1\text{H} + ({}^4\text{He}-X^-)$ (and to a lesser degree $({}^2\text{H}-X^-) + {}^4\text{He} \rightarrow {}^2\text{H} + ({}^4\text{He}-X^-)$).

The results of the present Monte-Carlo analysis are shown in Figs. 4 and 5. Shown are the likelihood areas in the Y_X - τ_X plane by different shading (coloring) that $> 99\%$, $95\% - 99\%$, $80\% - 95\%$, $20\% - 80\%$, $5\% - 20\%$, $1\% - 5\%$, and $< 1\%$ of all randomly generated models at the same Y_X and τ_X obey the observational constraints. Note that the CHAMP-to-entropy ratio Y_X is easily converted to Ω_X , the fractional contribution of the X -particle to the present critical density, if it had not decayed, via $\Omega_X h^2 = 2.73 \times 10^{11} Y_X (M_X/1 \text{ TeV})$, where h is the present Hubble constant in units $100 \text{ kms}^{-1}\text{Mpc}^{-1}$. Note also, that in all constraint figures Y_X denotes the total CHAMP-to-entropy ratio, assuming that only half of all CHAMPS are negatively charged. Whereas Fig. 4 shows results when the CHAMP decay is not associated with electromagnetic- or hadronic- energy release, Fig. 5 assumes a 100% electromagnetic CHAMP

decay and the associated electromagnetic cascade nucleosynthesis. It is evident from Fig. 4 that, when only effects of bound states are taken into account but not injection of energy during the decay, the probability distribution is extremely flat. In particular, whereas one still finds a small fraction of models which fail for Y_x as low as $\sim 10^{-19}$ only at $Y_x \gtrsim 10^{-12}$ one may exclude CHAMP BBN at the $> 99\%$ confidence level. Conservatively, and in the absence of more reliable rates, even Y_x as large as 10^{-12} may thus not reliably be ruled out. This is five orders of magnitude less stringent than the initial claim [2]. Note, however, that such large Y_x may only be (conservatively) acceptable for invisible, mass-degenerate, or stable CHAMPs, associated with very little (or no) injection of electromagnetically interacting energy (i.e. high-energy γ 's and $e^\pm$'s). This may be seen from Fig. 5, where for larger life times $Y_x \gtrsim 10^{-15}$ is ruled out at the $> 99\%$ level. Here models are ruled out principally due to violating the ${}^3\text{He}/{}^2\text{H}$ upper limit due to ${}^4\text{He}$ photodisintegration. This fact is essentially not changed by the X -particle being charged, as may be seen by inspection of the results in Ref. [14] for neutral decaying particles or by the dashed lines in Fig. 5, which shows the analogous limit for neutral particles. Immediately below the constraint line already between $20 - 80\%$ of all models yield acceptable abundance yields. This parameter space, is thus, currently not ruled out. However, it is conceivable that constraints on CHAMPs may be significantly tightened in future, to values possibly as low as $Y_X \lesssim 10^{-17} - 10^{-18}$, in case rates for the important reactions ${}^4\text{He}({}^2\text{H}-X^-, X^-){}^6\text{Li}$, $({}^1\text{H}-X^-) + {}^2\text{H} \rightarrow ({}^2\text{H}-X^-) + {}^1\text{H}$, and $({}^1\text{H}-X^-) + {}^4\text{He} \rightarrow ({}^4\text{He}-X^-)$ are determined, and contrive to yield unacceptably large ${}^6\text{Li}$ at low Y_X . Here it is noted that the by far most important ${}^6\text{Li}$ producing reaction, potentially leading to such stringent constraints, is ${}^4\text{He}({}^2\text{H}-X^-, X^-){}^6\text{Li}$ rather than ${}^2\text{H}({}^4\text{He}-X^-, X^-){}^6\text{Li}$, yielding the bulk of the ${}^6\text{Li}$ at temperatures $T \lesssim 1 \text{ keV}$ and not $T \approx 6 - 8 \text{ keV}$. Important here is the CHAMP exchange reaction $({}^1\text{H}-X^-) + {}^2\text{H} \rightarrow ({}^2\text{H}-X^-) + {}^1\text{H}$ which may continuously produce $({}^2\text{H}-X^-)$ bound states. (cf. Ref. [12] for more detail). This trend may be seen in Fig. 6, where ${}^6\text{Li}/{}^7\text{Li}$ isocontours are shown for a CHAMP with $B_h = 0$, and when the Born approximation for all rates is utilised. It is seen that for $\tau_x \gtrsim 10^6 \text{ sec}$ factor ~ 1000 more ${}^6\text{Li}$ is synthesised than for $\tau_x \lesssim 10^6 \text{ sec}$.

It is evident from Figs. 4 and 5 that the probability distributions only flattens significantly for CHAMP life times $\tau_x \gtrsim 5 \times 10^5 \text{ sec}$. For $10^5 \text{ sec} \lesssim \tau_x \lesssim 5 \times 10^5 \text{ sec}$ the probability passes from $> 99\%$ observationally acceptable models to $< 1\%$ observationally acceptable models within one decade of Y_x . This is because for shorter τ_x possible late-time destruction of ${}^6\text{Li}$ and ${}^7\text{Li}$ may not yet be efficient due to large $({}^1\text{H}-X^-)$ bound state fractions only forming at $T \lesssim 1 \text{ keV}$. Up to $\tau_x \lesssim 5 \times 10^5 \text{ sec}$ one may therefore approximately use the bounds on CHAMPs as given in Section 2. When $\tau_x \gtrsim 5 \times 10^5 \text{ sec}$ late-time processing (destruction) of ${}^6\text{Li}$, ${}^7\text{Li}$, and ${}^2\text{H}$ may occur ef-

ficiently. For long life times it is thus proposed to use *only* the constraint imposed by possible ${}^3\text{He}/{}^2\text{H}$ overproduction. This constraint is not affected by bound states and the uncertainties in reactions including bound states. The ${}^3\text{He}$ isotope is special in that it may not be destroyed without destroying ${}^2\text{H}$ as well. This is because the reaction ${}^3\text{He}({}^1\text{H}-X^-, X^-){}^4\text{Li}$ is endothermic and other reactions not involving ${}^2\text{H}$ nuclei are Coulomb suppressed. It is thus conservative to use the ${}^3\text{He}/{}^2\text{H}$ constraint, in particular, since there are other ${}^2\text{H}$ destroying (but not producing) reactions within bound state nucleosynthesis. This translates into using the constraints given in Ref. [14] for $B_h = 0$ (and shown by the dashed curve for $\tau_x \gtrsim 10^6 \text{ sec}$ in Fig. 5). Since the fraction of rest mass f_{EM} which is converted to electromagnetic interacting energy has been assumed $f_{EM} = 1$ for $B_h = 0$ in the figures of Ref. [14], the constraint has to be rescaled accordingly, when neutrino losses are significant, or close-to mass degeneracy between mother and daughter particle exists. Finally, in the window $5 \times 10^5 \text{ sec} \lesssim \tau_x \lesssim 3 \times 10^6 \text{ sec}$ one may apply an additional constraint, stronger than that due to ${}^3\text{He}/{}^2\text{H}$ overproduction. From Figs. 4 and 5 it is found that for $Y_x \gtrsim 3 \times 10^{-14} - 10^{-13}$ only $\lesssim 1\%$ of all models are observationally acceptable.

A procedure very similar to this has been very recently applied by Ref. [18] to derive a lower limit on the gaugino mass parameter $m_{1/2}$ in the constrained minimal supersymmetric standard model (CMSSM) when the gravitino is the lightest supersymmetric particle (LSP). This study considered only the catalyzed reaction ${}^2\text{H} + ({}^4\text{He}-X^-) \rightarrow {}^6\text{Li} + X^-$ for intermediate life times of the next-to-LSP (NLSP), arguing that even without the ${}^6\text{Li}$ bound long NLSP life times were already ruled out priorly by the ${}^3\text{He}/{}^2\text{H}$ upper limit. Since in the parameter space under investigation the NLSP is the stau, which has small hadronic branching ratio, the conclusion for $\tau_x \lesssim 5 \times 10^5 \text{ sec}$ is thus not expected to change much when the present results are used. Using the results of the Monte-Carlo analysis shown in Figs. 4 and 5 typical stau-to-entropy ratios $Y_{\tilde{\tau}} \gtrsim 10^{-13}$ should be also disallowed for $5 \times 10^5 \text{ sec} \lesssim \tau_x \lesssim 3 \times 10^6 \text{ sec}$ due to ${}^6\text{Li}$ overproduction and for $\tau_x \gtrsim 3 \times 10^6 \text{ sec}$ due to ${}^3\text{He}/{}^2\text{H}$ overproduction, rendering the conclusions of Ref. [18] likely unchanged.

IV. CONCLUSIONS

In this letter the results of a detailed study of constraints on charged massive particles X^- from Big Bang nucleosynthesis was presented. It was pointed out earlier that bound states between ${}^4\text{He}$ and negatively charged X^- may lead to the efficient catalytic production of ${}^6\text{Li}$ at $T \approx 6 - 8 \text{ keV}$ [2]. Recently I have shown [12], that BBN with CHAMPs enters a second phase of nucleosynthesis at $T \lesssim 1 \text{ keV}$, capable of destroying all priorly synthesized ${}^6\text{Li}$. Altogether nineteen reactions important for late-time BBN were identified. When these processes are included in the analysis, drastic changes concerning lim-

its on the existence of CHAMPs when compared to those priorly derived [2, 7, 9, 10, 11], are obtained.

Limits on the existence of CHAMPs during and after BBN are derived in two different decay time ranges. For very short life times $\tau_x \lesssim 3 \times 10^2 \text{sec}$ limits are independent of the decaying particle being charged, or not. In the range $3 \times 10^2 \text{sec} \lesssim \tau_x \lesssim 5 \times 10^5 \text{sec}$ the important rates (i.e. ${}^2\text{H}({}^4\text{He}-X^-, X^-){}^6\text{Li}$) are relatively well determined [9], such that a Monte-Carlo analysis may be avoided. It is stressed that the approximation of the ${}^4\text{He}$ -bound state fraction by the Saha equation leads to a factor ~ 10 overestimate in the synthesized ${}^6\text{Li}$ abundance. Due to substantial reaction rate uncertainties a full Monte-Carlo analysis had to be performed to obtain reliable and conservative bounds in the decay time range $5 \times 10^5 \text{sec} \lesssim \tau_x \lesssim 10^{12} \text{sec}$. It was found that when a number of reaction rates are large, and when electromagnetic energy injection is absent, baryon-to-entropy ratios Y_x as large as $\gtrsim 10^{-12}$ may be observationally acceptable. On the other hand, in case a number of reaction rates are determined more precisely, in particular the rates for ${}^4\text{He}({}^2\text{H}-X^-, X^-){}^6\text{Li}$, $({}^1\text{H}-X^-) + {}^2\text{H} \rightarrow ({}^2\text{H}-X^-) + {}^1\text{H}$, and $({}^1\text{H}-X^-) + {}^4\text{He} \rightarrow ({}^1\text{H}-X^-) + {}^4\text{He}$, it may be conceivable that limits on long-lived CHAMPs are improved by orders of magnitude, with the CHAMP-to-entropy ratio possibly constrained to be below $10^{-17} - 10^{-18}$. A prescription is given for how to place conservative limits on CHAMPs, given current reaction rate uncertainties.

The final result of this study is somewhat surprising. When electromagnetic- and hadronic- energy release are included, and within the reaction rate uncertainties, conservative limits on charged decaying particles are no stronger than those on neutral particles. The exception here is the decay time range $10^3 - 3 \times 10^6 \text{sec}$ but only when the hadronic branching ratio is small $B_h \lesssim 10^{-2} - 10^{-1}$, as is the case for supersymmetric staus.

I acknowledge helpful discussions with and M. Asplund, S. Bailly, O. Kartavtsev, K. Kohri, A. Korn, G. Moulta, M. Pospelov, J. Rafelski, G. Starkman, F. Steffen, V. Tatischeff, and T. Yanagida.

V. ERRATUM: BOUNDS ON LONG-LIVED CHARGED MASSIVE PARTICLES FROM BIG BANG NUCLEOSYNTHESIS

A recent detailed computation by Kamimura *et al.*[20] of the most important reaction rates entering bound state nucleosynthesis (see also the discussion in Ref. [21]) has established that late time $\tau \gtrsim 10^6 \text{sec}$ processing of light element abundances as envisioned possible by the present author [12] does not usually take place. This is mostly because of the abundance of proton- X^- bound states staying small due to efficient exchange reactions transfer-

ring X^- from p to ${}^4\text{He}$, but also due to Coulomb barriers between p and nuclei being only partially shielded when protons are in bound states. The Monte Carlo analysis as




FIG. 7: Limits on the primordial CHAMP-to-entropy ratio $Y_x = n_X/s$ (with $n_X/s = Y_x/2$) as a function of their life time τ_x . Shown are constraint lines for CHAMPs of mass $M_x = 1 \text{TeV}$ and a variety of hadronic branching ratios $B_h = 10^{-5} - 1$, as labeled in the figure. Solid (red) lines correspond to the conservative limit ${}^6\text{Li}/{}^7\text{Li} < 0.66$, whereas dashed (blue) lines correspond to ${}^6\text{Li}/{}^7\text{Li} < 0.1$. It is seen that only for CHAMPs with $B_h \lesssim 10^{-2}$ the effects of bound states become important. For small decay times $\tau_x \lesssim 10^3 \text{sec}$ and large decay times $\tau_x \gtrsim 10^7 \text{sec}$ the limits on CHAMP abundances are virtually identical to those on the abundance of neutral relic decaying particles [14].




FIG. 8: As Fig. 7 but for $M_x = 100 \text{GeV}$.

performed in Section II and III is therefore superfluous. Figs. 4-6 of the paper are replaced by Figs. 7 and 8 in this erratum. It is cautioned, however, that for relatively large $Y_X > 2Y_{\text{He}}$ late time processing may still occur to some degree.

-
- [1] A. De Rujula, S. L. Glashow and U. Sarid, Nucl. Phys. B **333**, 173 (1990); S. Dimopoulos, D. Eichler, R. Esmailzadeh and G. D. Starkman, Phys. Rev. D **41**, 2388 (1990); J. Rafelski, M. Sawicki, M. Gajda and D. Harley, Phys. Rev. A **44**, 4345 (1991).
 - [2] M. Pospelov, arXiv:hep-ph/0605215.
 - [3] K. Kohri and F. Takayama, arXiv:hep-ph/0605243.
 - [4] M. Kaplinghat and A. Rajaraman, Phys. Rev. D **74**, 103004 (2006).
 - [5] C. Bird, K. Koopmans and M. Pospelov, arXiv:hep-ph/0703096.
 - [6] T. Jittoh, K. Kohri, M. Koike, J. Sato, T. Shimomura and M. Yamanaka, arXiv:0704.2914 [hep-ph].
 - [7] R. H. Cyburt, J. R. Ellis, B. D. Fields, K. A. Olive and V. C. Spanos, JCAP **0611**, 014 (2006).
 - [8] V. V. Smith, D. L. Lambert, and P. E. Nissen, Astrophys. J. **408**, 262 (1993); **506**, 405 (1998); L. M. Hobbs and J. A. Thorburn, Astrophys. J. **491**, 772 (1997); R. Cayrel, M. Spite, F. Spite, E. Vangioni-Flam, M. Cassé, and J. Audouze, Astron. & Astrophys. **343**, 923 (1999); P. E. Nissen, M. Asplund, V. Hill, and S. D'Odorico, Astr. & Astrophys. **357**, L49 (2000); M. Asplund, D. L. Lambert, P. E. Nissen, F. Primas and V. V. Smith, Astrophys. J. **644**, 229 (2006).
 - [9] K. Hamaguchi, T. Hatsuda, M. Kamimura, Y. Kino and T. T. Yanagida, arXiv:hep-ph/0702274.
 - [10] M. Kawasaki, K. Kohri and T. Moroi, Phys. Lett. B **649**, 436 (2007).
 - [11] J. Pradler and F. D. Steffen, Phys. Lett. B **648**, 224 (2007).
 - [12] K. Jedamzik, arXiv:0707.2070 [astro-ph].
 - [13] The effects of annihilation of X^- with X^+ in $X^- - X^+$ bound states forming to some degree [5] are subdominant compared to those of the decay itself.
 - [14] K. Jedamzik, Phys. Rev. D **74**, 103509 (2006).
 - [15] A. J. Korn *et al.*, Nature **442**, 657 (2006).
 - [16] M. H. Pinsonneault, G. Steigman, T. P. Walker, . K. Narayanans and V. K. Narayanan, Astrophys. J. **574**, 398 (2002); O. Richard, G. Michaud, and J. Richer, Astrophys. J., **619**, 538 (2005).
 - [17] S. Dimopoulos, R. Esmailzadeh, L. J. Hall and G. D. Starkman, Astrophys. J. **330**, 545 (1988).
 - [18] J. Pradler and F. D. Steffen, arXiv:0710.2213 [hep-ph].
 - [19] F. Takayama, arXiv:0704.2785 [hep-ph].
 - [20] M. Kamimura, Y. Kino and E. Hiyama, arXiv:0809.4772 [nucl-th].
 - [21] M. Pospelov, J. Pradler and F. D. Steffen, JCAP **0811**, 020 (2008).